



Endari Prior Authorization Program Summary

This program applies to Medicaid.

The BCBS MN Step Therapy Supplement applies to this program for Medicaid.

Requests for an oral liquid form of a drug must be approved if BOTH of the following apply:

- 1) the indication is FDA approved AND
- 2) the patient is using an enteral tube for feeding or medication administration

POLICY REVIEW CYCLE

Effective Date
03-01-2024

Date of Origin
04-01-2018

FDA APPROVED INDICATIONS AND DOSAGE

Agent(s)	FDA Indication(s)	Notes	Ref#
Endari® (L-glutamine) Oral powder	To reduce the acute complications of sickle cell disease in adult and pediatric patients 5 years of age and older		

See package insert for FDA prescribing information: <https://dailymed.nlm.nih.gov/dailymed/index.cfm>

CLINICAL RATIONALE

Sickle Cell Disease	<p>Sickle cell disease (SCD) is the name given to a group of lifelong inherited conditions that affect hemoglobin. People with SCD have atypical hemoglobin molecules called hemoglobin S, which can distort red blood cells into a sickle or crescent shape.(2)</p> <p>Signs and symptoms of SCD usually begin in early childhood. Characteristic features of SCD include anemia, repeated infections, and periodic episodes of pain. The severity of symptoms varies from person to person and can range from mild to requiring frequent hospitalizations.(2)</p> <p>SCD effects nearly every system in the body. SCD has both acute and chronic complications. An episode of severe pain [acute vaso-occlusive crisis (VOC)] is the most common acute complication of SCD. In addition to VOCs other common acute complications of SCD include fever related to infection, acute kidney injury (AKI), hepatobiliary complications, acute anemia, splenic sequestration, acute chest syndrome (ACS), and acute stroke. Chronic complications of SCD can affect almost any organ, and certain acute complications often evolve into chronic phases. The most common chronic complications of SCD include chronic pain, chronic anemia, avascular necrosis, leg ulcers, pulmonary hypertension, renal complications, stuttering/recurrent priapism, and ophthalmologic complications.(2)</p>
---------------------	--

Pain is the most common complication of SCD for both acute and chronic complications. Pain can be acute, chronic, or an acute episode superimposed on chronic pain. In SCD, pain is considered chronic if it lasts more than 3 months. People with SCD experience both nociceptive and neuropathic pain.(2)

Recurrent and unpredictable episodes of vaso-occlusion are the hallmark of sickle cell disease. Discoveries over the past 2 decades have highlighted the important contributions of various cellular and soluble participants in the vaso-occlusive cascade. Although the molecular basis of SCD is well characterized, the complex mechanisms underlying VOC have not been fully elucidated. Based on direct observations in SCD mice, adhesive interactions of SS-RBCs and leukocytes to the endothelium play important roles in the initiation of VOC. It is thought that the activated adherent leukocytes, which are rigid and larger than sickle cell-red blood cells (SS-RBC), likely drive VOC in collecting venules, whereas the SS-RBCs may contribute in smaller vessels or in situations where there is no potent inflammatory trigger.(4)

Triggers for VOC vary and can include inflammation, stress, increased viscosity, decreased flow, hemolysis, or a combination of the following factors:(4)

- Endothelial activation by SS-RBCs and other inflammatory mediators
- Recruitment of adherent leukocytes
- Activation of recruited neutrophils and of other leukocytes (e.g., monocytes or iNKT cells)
- Interactions of sickle erythrocytes with adherent neutrophils
- Vascular clogging by heterotypic cell-cell aggregates composed of SS-RBCs, adherent leukocytes and possibly platelets
- Increased transit time to greater than the delay time for deoxygenation-induced hemoglobin polymerization, propagating retrograde VOC
- Ischemia as a result of the obstruction that creates a feedback loop of worsening endothelial activation

Sickle hemoglobin can cause damage to the RBC membrane from deformation by polymer formation. In addition, the mutated globin can undergo autooxidation and precipitate on the inner surface of the RBC membrane, causing membrane damage via iron-mediated generation of oxidants. Both endothelial selectins, P-selectin and E-selectin, have been suggested to participate in VOC.(4)

Nearly all people with SCD have chronic anemia, but individual baseline hemoglobin values vary widely depending upon hemoglobin genotype (HbSS, HbSC, HbS β^+ -thalassemia, HbS β^0 -thalassemia). It is important for the patient and the primary care provider to know the baseline or "steady state" hemoglobin value to inform ongoing monitoring and management during acute complications.(2)

Hydroxyurea, a ribonucleotide reductase inhibitor, was identified as an option to increase fetal hemoglobin (HbF) levels in people with SCD. The initial clinical trial of hydroxyurea for the treatment of sickle cell anemia (SCA) involved two people. The results of this study showed favorable outcomes which lead to two extended studies with larger cohorts of people. Although HbF induction is the most powerful effect of hydroxyurea and provides the biggest direct benefit for people who have SCD,

	<p>additional mechanisms of actions and benefits exist. Hydroxyurea lowers the number of circulating leukocytes and reticulocytes and alters the expression of adhesion molecules, all of which contribute to vaso-occlusion. Hydroxyurea also raises RBC volume [higher mean corpuscular volume (MCV)] and improves cellular deformability and rheology, which increases blood flow and reduces vaso-occlusion.(3)</p> <p>An expert panel report of evidence-based management of sickle cell disease supports the use of hydroxyurea with strong recommendations in the following:(3)</p> <ul style="list-style-type: none"> • In adults with SCA who have three or more sickle cell-associated moderate to severe pain crises in a 12 month period • In adults with SCA who have sickle cell-associated pain that interferes with daily activity and quality of life • In adults with SCA who have a history of severe and/or recurrent acute coronary syndrome (ACS) • In adults with SCA who have severe symptomatic chronic anemia that interferes with daily activities or quality of life • In infants 9 months of age and older, children, and adolescents with SCA, offer treatment with hydroxyurea regardless of clinical severity to reduce SCD-related complications (e.g., pain, dactylitis, ACS, anemia) <p>A clinical response to treatment with hydroxyurea may take 3-6 months. Therefore, the expert panel report of evidence-based management of sickle cell disease recommends a 6 month trial on the maximum tolerated dose is required prior to considering discontinuation due to treatment failure, whether due to lack of adherence or failure to respond to therapy.(3)</p>
Efficacy (1)	<p>The mechanism of action of the amino acid L-glutamine in treating sickle cell disease (SCD) is not fully understood. Oxidative stress phenomena are involved in the pathophysiology of SCD. Sickle red blood cells (RBCs) are more susceptible to oxidative damage than normal RBCs, which may contribute to the chronic hemolysis and vaso-occlusive events associated with SCD. The pyridine nucleotides, NAD⁺ (oxidized nicotinamide adenine dinucleotide) and its reduced form NADH (nicotinamide adenine dinucleotide + hydrogen), play roles in regulating and preventing oxidative damage in RBCs. L-glutamine may improve the NAD redox potential in sickle RBCs through increasing the availability of reduced glutathione.</p> <p>The efficacy of L-glutamine was evaluated in a randomized, double-blind, placebo controlled, multi-center clinical trial with 230 patients. Efficacy was demonstrated by a reduction in the number of sickle cell crises through Week 48 and prior to the start of tapering among patients that received L-glutamine compared to patients who received placebo. The recurrent crisis event time analysis yielded an intensity rate ratio (IRR) value of 0.75 with 95% CI= (0.62, 0.90) and (0.55, 1.01) based on unstratified models using the Andersen-Gill and Lin, Wei, Yang and Ying methods, respectively in favor of L-glutamine, suggesting that over the entire 48-week period, the average cumulative crisis count was reduced by 25% from the L-glutamine group over the placebo group.</p>
Safety (1)	Endari (L-glutamine) has no FDA labeled contraindications.

REFERENCES

Number	Reference
1	Endari prescribing information. Emmaus Medical, Inc. October 2020.

Number	Reference
2	U.S. National Library of Medicine. Genetics Home Reference. Sickle cell disease. November 2019.
3	U.S. Department of Health and Human Services. National Institute of Health. Evidence-Based Management of Sickle Cell Disease. Expert Panel Report, 2014.
4	Manwani D, Frenette PS, Vaso-occlusion in sickle cell disease: pathophysiology and novel targeted therapies. Blood. 2013 Dec 5; 122(24): 3892-3898.

POLICY AGENT SUMMARY PRIOR AUTHORIZATION

Target Brand Agent(s)	Target Generic Agent(s)	Strength	Targeted MSC	Available MSC	Final Age Limit	Preferred Status
Endari	glutamine (sickle cell) powd pack	5 GM	M ; N ; O ; Y	N		

CLIENT SUMMARY – PRIOR AUTHORIZATION

Target Brand Agent Name(s)	Target Generic Agent Name(s)	Strength	Client Formulary
Endari	glutamine (sickle cell) powd pack	5 GM	

PRIOR AUTHORIZATION CLINICAL CRITERIA FOR APPROVAL

Module	Clinical Criteria for Approval
	<p>Initial Evaluation</p> <p>Target Agent(s) will be approved when ONE of the following is met:</p> <ol style="list-style-type: none"> 1. ALL of the following <ol style="list-style-type: none"> A. The patient has a diagnosis of sickle cell disease AND B. The patient is using the requested agent to reduce the acute complications of sickle cell disease AND C. If the patient has an FDA approved indication, then ONE of the following: <ol style="list-style-type: none"> 1. The patient’s age is within FDA labeling for the requested indication for the requested agent OR 2. The prescriber has provided information in support of using the requested agent for the patient’s age AND D. ONE of the following <ol style="list-style-type: none"> 1. The patient’s medication history includes hydroxyurea AND ONE of the following: <ol style="list-style-type: none"> A. The patient has had an inadequate response to hydroxyurea OR B. The prescriber has submitted an evidence-based and peer-reviewed clinical practice guideline supporting the use of the requested agent over hydroxyurea OR 2. The patient has an intolerance or hypersensitivity to hydroxyurea OR 3. The patient has an FDA labeled contraindication to hydroxyurea OR 4. The patient is currently being treated with the requested agent as indicated by ALL of the following: <ol style="list-style-type: none"> A. A statement by the prescriber that the patient is currently taking the requested agent AND B. A statement by the prescriber that the patient is currently receiving a positive therapeutic outcome on requested agent AND

Module	Clinical Criteria for Approval
	<p style="text-align: center;">C. The prescriber states that a change in therapy is expected to be ineffective or cause harm OR</p> <p>5. The prescriber has provided documentation that hydroxyurea cannot be used due to a documented medical condition or comorbid condition that is likely to cause an adverse reaction, decrease ability of the patient to achieve or maintain reasonable functional ability in performing daily activities or cause physical or mental harm AND</p> <p>E. ONE of the following:</p> <ol style="list-style-type: none"> 1. The patient will NOT be using the requested agent in combination with Adakevo (crizanlizumab-tmca) OR Oxbryta (voxelotor) OR 2. Information has been provided supporting the use of the requested agent in combination with Adakevo (crizanlizumab-tmca) or Oxbryta (voxelotor) AND <p>F. The patient does NOT have any FDA labeled contraindications to the requested agent AND</p> <p>G. The requested quantity (dose) does NOT exceed the maximum FDA labeled dose for the requested indication OR</p> <p>2. If the request is for an oral liquid form of medication, then BOTH of the following:</p> <ol style="list-style-type: none"> A. The patient has an FDA approved indication AND B. The patient uses an enteral tube for feeding or medication administration <p>Length of Initial Approval: 12 months</p> <p>Renewal Evaluation</p> <p>Target Agent(s) will be approved when ONE of the following is met:</p> <ol style="list-style-type: none"> 1. ALL of the following: <ol style="list-style-type: none"> A. The patient has been previously approved through the plan's Prior Authorization process AND B. The patient has had clinical benefit with the requested agent (i.e., reduction in acute complications of sickle cell disease since initiating therapy with the requested agent) AND C. ONE of the following: <ol style="list-style-type: none"> 1. The patient will NOT be using the requested agent in combination with Adakevo (crizanlizumab-tmca) OR Oxbryta (voxelotor) OR 2. Information has been provided supporting the use of the requested agent in combination with Adakevo (crizanlizumab-tmca) or Oxbryta (voxelotor) AND D. The patient does NOT have any FDA labeled contraindications to the requested agent AND E. The requested quantity (dose) does NOT exceed the maximum FDA labeled dose for the requested indication OR 2. If the request is for an oral liquid form of medication, then BOTH of the following: <ol style="list-style-type: none"> A. The patient has an FDA approved indication AND B. The patient uses an enteral tube for feeding or medication administration <p>Length of Renewal Approval: 12 months</p>

