# BULLETIN

Blue Cross and Blue Shield of Minnesota and Blue Plus

June 12, 2008

# July 2008 HCPCS code update

Effective July 1, 2008, Blue Cross and Blue Shield of Minnesota and Blue Plus will accept the quarterly Level II (alphanumeric) and Level I (CPT) HCPCS\* additions, revisions and deletions published by the Centers of Medicare & Medicaid Services (CMS) and the American Medical Association (AMA).

### Submitting codes

Blue Cross will accept all added HCPCS codes effective on July 1, 2008.

#### Code list

HCPCS and CPT manuals are generally only published once a year; however, CMS or AMA may publish added, revised and/or deleted codes on a quarterly basis. The codes listed below are effective July 1, 2008 unless otherwise specified.

#### ADDED CODES:

CG Policy criteria applied  C9242 Injection, fosaprepitant, 1 mg  C9356 Tendon, porous matrix of cross-linked collagen and glycosaminoglycan matrix (TenoGlide Tendon Protector Sheet), per square centimeter  C9357 Dermal substitute, granulated cross-linked collagen and glycosaminoglycan matrix (Flowable Wound Matrix), 1 cc  C9358 Dermal substitute, native, non-denatured collagen (SurgiMend Collagen Matrix), per 0.5 square centimeters  G0398 Home sleep study test (HST) with type II portable monitor, unattended; minimum of 7 channels: EEG, EOG, EMG, ECG/heart reate, airflow, respiratory effort and oxygen saturation  G0399 Home sleep test (HST) with type II portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation  G0400 Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  90738 Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  3351F No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  3353F Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  3354F Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to ode for primary service)  10190T Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  10191T Insertion of anterior segment aqueous drainage device, without extraocular reservoir; external a		
Tendon, porous matrix of cross-linked collagen and glycosaminoglycan matrix (TenoGlide Tendon Protector Sheet), per square centimeter  Dermal substitute, granulated cross-linked collagen and glycosaminoglycan matrix (Flowable Wound Matrix), 1 cc  Dermal substitute, native, non-denatured collagen (SurgiMend Collagen Matrix), per 0.5 square centimeters  Home sleep study test (HST) with type II portable monitor, unattended; minimum of 7 channels: EEG, EOG, EMG, ECG/heart reate, airflow, respiratory effort and oxygen saturation  Home sleep test (HST) with type III portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation  Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to primary procedure)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)	CG	Policy criteria applied
square centimeter  C9357 Dermal substitute, granulated cross-linked collagen and glycosaminoglycan matrix (Flowable Wound Matrix), 1 cc  C9358 Dermal substitute, native, non-denatured collagen (SurgiMend Collagen Matrix), per 0.5 square centimeters  G0398 Home sleep study test (HST) with type II portable monitor, unattended; minimum of 7 channels: EEG, EOG, EMG, ECG/heart reate, airflow, respiratory effort and oxygen saturation  G0399 Home sleep test (HST) with type III portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation  G0400 Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  90738 Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  3352F No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  3353F Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	C9242	Injection, fosaprepitant, 1 mg
Dermal substitute, native, non-denatured collagen (SurgiMend Collagen Matrix), per 0.5 square centimeters  G0398 Home sleep study test (HST) with type II portable monitor, unattended; minimum of 7 channels: EEG, EOG, EMG, ECG/heart reate, airflow, respiratory effort and oxygen saturation  G0399 Home sleep test (HST) with type III portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation  G0400 Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  S353F Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	C9356	
Home sleep study test (HST) with type II portable monitor, unattended; minimum of 7 channels: EEG, EOG, EMG, ECG/heart reate, airflow, respiratory effort and oxygen saturation  Home sleep test (HST) with type III portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation  Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	C9357	Dermal substitute, granulated cross-linked collagen and glycosaminoglycan matrix (Flowable Wound Matrix), 1 cc
ECG/heart reate, airflow, respiratory effort and oxygen saturation  Home sleep test (HST) with type III portable monitor, unattended; minimum of 4 channels: 2 respiratory movement/airflow, 1 ECG/heart rate and 1 oxygen saturation  Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	C9358	Dermal substitute, native, non-denatured collagen (SurgiMend Collagen Matrix), per 0.5 square centimeters
Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels  Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	G0398	
Japanese encephalitis virus vaccine, inactivated, for intramuscular use  Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	G0399	
Negative screen for depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	G0400	Home sleep test (HST) with type IV portable monitor, unattended; minimum of 3 channels
(MDD)  No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	90738	Japanese encephalitis virus vaccine, inactivated, for intramuscular use
Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	3351F	
Clinically significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	3352F	No significant depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)
(MDD)  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	3353F	Mild to moderate depressive symptoms as categorized by using a standardized depression screening/assessment too (MDD)
injured patient; first 30-74 minutes  Remote real-time interactive videoconferenced critical care, evaluation and management of the critically ill or critically injured patient; each additional 30 minutes (List separately in addition to code for primary service)  Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	3354F	
injured patient; each additional 30 minutes (List separately in addition to code for primary service)  O190T Placement of intraocular radiation source applicator (List separately in addition to primary procedure)  O191T Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	0188T	
0191T Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach	0189T	
	0190T	Placement of intraocular radiation source applicator (List separately in addition to primary procedure)
0192T Insertion of anterior segment aqueous drainage device, without extraocular reservoir; external approach	0191T	Insertion of anterior segment aqueous drainage device, without extraocular reservoir; internal approach
	0192T	Insertion of anterior segment aqueous drainage device, without extraocular reservoir; external approach



## **REVISED CODE:**

0124T	Conjunctival incision with posterior juxtaseleral extraseleral placement of pharmacological agent (does not include supply of
	medication)

# **DISCONTINUED CODES** (the following codes were discontinued effective 12/31/2007):

	, 0
G0297	Insertion of single chamber pacing cardioverter defibrillator pulse generator
G0377	Administration of vaccine for Part D drug

#### Questions?

If you have any questions, please contact provider service at (651) 662-5200 or 1-800-262-0820.

CPT codes copyright 2008 American Medical Association. All Rights Reserved. CPT is a trademark of the AMA.

\*HCPCS stands for Health Care Procedure Coding System

